

A NOTE ON THE DEFORMED HERMITIAN YANG-MILLS PDE

VAMSI P. PINGALI

ABSTRACT. We prove *a priori* estimates for a generalised Monge-Ampère PDE with “non-constant coefficients” thus improving a result of Sun in the Kähler case. We apply this result to the deformed Hermitian Yang-Mills (dHYM) equation of Jacob-Yau to obtain an existence result and *a priori* estimates for some ranges of the phase angle assuming the existence of a subsolution. We then generalise a theorem of Collins-Székelyhidi on toric varieties and use it to address a conjecture of Collins-Jacob-Yau.

1. INTRODUCTION

According to some versions of superstring theory the spacetime of the universe is constrained to be a product (more generally a fibration) of a compact Calabi-Yau three-fold and a four dimensional Lorentzian manifold. A “duality” relates the geometry of this Calabi-Yau manifold with another “mirror” Calabi-Yau manifold. From a differential geometry standpoint this maybe thought of (roughly) as a relationship between the existence of “nice” metrics on a line bundle on one Calabi-Yau manifold and special Lagrangian submanifolds of the other Calabi-Yau manifold. Using the Fourier-Mukai transform on a torus fibration, Leung-Yau-Zaslow suggested [6] that this “mirror symmetry” implies that the following special Lagrangian-type PDE ought to be satisfied on a compact Kähler manifold (X, ω) of complex dimension n .

$$(1.1) \quad \operatorname{Im}((\omega - F)^n) = \tan(\hat{\theta}) \operatorname{Re}((\omega - F)^n),$$

where F is the curvature of the Chern connection of a hermitian metric on a holomorphic line bundle L and $\tan(\theta)$ is the constant defined by integrating equation 1.1 on both sides. The angle $\hat{\theta}$ has been christened “the phase angle”.

In [5], Jacob and Yau showed that for a given ample line bundle L over a compact Kähler manifold with non-negative orthogonal bisectional curvature, L^k admits a solution to the equation 1.1. In [1] Collins, Jacob and Yau showed existence under the assumption of a subsolution with supercritical phase. In this paper we aim to get similar results using a different method.

We first prove *a priori* estimates for a generalised Monge-Ampère PDE with “non-constant” coefficients. In the Kähler case this is an improvement of a result of Sun [9].

Theorem 1.1. Assume that $n\Omega^{n-1} - \sum_{k=0}^{n-1} \binom{n}{k} c_k(n-k)\Omega^{n-k-1}\omega^k > 0$. Let ϕ satisfying $\inf_X \phi = 1$ be a smooth solution of

$$(1.2) \quad \Omega_\phi^n = \sum_{k=0}^{n-1} \binom{n}{k} c_k \Omega_\phi^{n-k} \omega^k,$$

where $c_k \geq 0$ are smooth functions such that either $c_k = 0$ or $c_k > 0$ throughout, and $\sum_k c_k > 0$. Then $\|\phi\|_{C^{2,\alpha}} \leq C(c_k, \Omega, \omega, X, \alpha)$.

We then apply the preceding result to equation 1.1 to arrive at the following result.

Theorem 1.2. Let (L, h_0) be a hermitian holomorphic line bundle on a compact Kähler n -complex dimensional manifold (X, ω) . Let F_0 denote the curvature of the Chern connection of (L, h_0) . Let $\hat{\theta}$ be a smooth function on X . Under the following assumptions any smooth solution $h = h_0 e^{-\hat{\phi}}$ of equation 1.1 satisfies the a priori estimate $\|\phi\|_{C^{2,\alpha}} \leq C(\alpha, X, \Omega, \hat{\theta}, \omega)$. In addition, if $\hat{\theta}$ is a constant such that the integrals on both sides of equation 1.1 are equal, then there exists a smooth solution satisfying 1.1.

- (1) If $n = 2m + 1$ then $\tan(\hat{\theta})$ is assumed to be smooth and $\Omega = \sqrt{-1}F_0 - \omega \tan(\hat{\theta})$ is assumed to be a supercritical phase subsolution, i.e., $\Omega > 0$ and $n\Omega^{n-1} - \sum c_k k \Omega^{k-1} \omega^{n-k} > 0$ where c_k are defined as

$$(1.3) \quad \begin{aligned} c_{2j} &= (-1)^{m+1+j} \binom{2m+1}{2j} \sec(\hat{\theta})^{2m+2-2j} \sin((2m-2j)\hat{\theta}) \\ c_{2j+1} &= (-1)^{m+j+1} \binom{2m+1}{2j+1} \sec(\hat{\theta})^{2m-2j+1} \cos((2m-2j-1)\hat{\theta}). \end{aligned}$$

In addition, we assume that c_k as defined above are either strictly positive throughout or are identically zero.

- (2) If $n = 2m$ then $\cot(\hat{\theta})$ is assumed to be smooth and $\Omega = \sqrt{-1}F_0 + \omega \cot(\hat{\theta})$ is assumed to be a supercritical phase subsolution, i.e., $\Omega > 0$ and $n\Omega^{n-1} - \sum c_k k \Omega^{k-1} \omega^{n-k} > 0$ where c_k are defined as

$$(1.4) \quad c_k = \csc(\hat{\theta})^{2m-k} (-1)^{2m-k+1} \binom{2m}{k} \sin((2m-k-1)\hat{\theta}).$$

In addition, we assume that c_k as defined above are either strictly positive throughout or are identically zero.

Remark 1.3. The assumption $\Omega > 0$ in theorem 1.2 is implied by the assumption of supercritical phase ($\sum \arctan(\mu_l) = \theta \in ((n-2)\frac{\pi}{2}, n\frac{\pi}{2})$) where μ_l are the eigenvalues of $\sqrt{-1}F$ with respect to ω) using proposition 8.1 and lemma 8.2 of [1].

Remark 1.4. Theorem 1.2 is actually a special case of theorems 1.1 and 1.2 of [1]. But the point here is not so much the theorem itself as the proof of it. The estimates involved and the method of continuity used are much simpler compared to those in [1].

The main assumption of having a subsolution in theorem 1.2 is actually quite difficult to check. It seems that it is almost as difficult as actually solving the equation itself. Therefore, it is desirable to have easier “algebro-geometric” assumptions on a generalised Monge-Ampère equation akin to the J-flow equation in [2]. Indeed, by modifying the technique in [2] we get the following result on toric varieties. Notice that the following result is also an improvement of [2] even in spirit because we are allowing $c_1 = 0$ in this theorem (at the cost of assuming that $c_n > 0$).

Theorem 1.5. Let X be a compact toric manifold of dimension n , with two toric Kähler metrics Ω and ω . Suppose that

$$\int_X \Omega^n - \sum_{k=1}^n \int_X \binom{n}{k} c_k \omega^k \Omega^{n-k} \geq 0,$$

where $c_k \geq 0$ are constants such that either $c_1 > 0$ or $c_n > 0$. Also assume that

$$\int_V \Omega^p - \sum_k \int_V c_k \binom{p}{k} \omega^k \Omega^{p-k} > 0$$

for all p -dimensional toric subvarieties and all $p \leq n - 1$. Then there is a smooth Kähler metric $\Omega_\phi = \Omega + \sqrt{-1}\partial\bar{\partial}\phi$ such that

$$(1.5) \quad \Omega_\phi^n = \sum_{k=1}^n c_k \binom{n}{k} \omega^k \Omega_\phi^{n-k} + d\omega^n$$

for a suitable constant $d \geq 0$ with equality holding if and only if equality holds in inequality 1.5.

Using this theorem we get the following corollary which addresses conjecture 1.4 of [1].

Corollary 1.6. *For every toric subvariety V of a toric manifold X , define $\theta_V = \text{Arg} \int_V (\omega - F)^{\dim V}$. Assume that ω and F_0 are torus invariant. Let $\hat{\theta}$ as in equation 1.1 be a constant. If*

- (1) *For every toric subvariety of X , $\Theta_V > \hat{\theta} - (n - \dim V)\frac{\pi}{2}$, and*
- (2) *c_k as defined in theorem 1.2 satisfy $c_k \geq 0$ with $c_0 > 0$,*

then there exists a smooth solution to deformed Hermitian Yang-Mills equation 1.1.

Almost all the results in this paper are based on the observation that equation 1.1 can be treated as a generalised Monge-Ampère equation. We believe that one can push these techniques further in order to obtain more general results. This will be investigated in future work.

Acknowledgements : The author thanks Gábor Székelyhidi for helpful discussions.

2. SOME *a priori* ESTIMATES FOR THE GENERALISED MONGE-AMPÈRE EQUATION

In this section we prove a fairly general *a priori* estimate on compact Kähler n -dimensional manifolds (X, ω) with another Kähler metric $\Omega_\phi = \Omega + \sqrt{-1}\partial\bar{\partial}\phi$. First one has a uniform estimate.

Lemma 2.1. *Assume that $n\Omega^{n-1} - \sum_{k=0}^{n-1} \binom{n}{k} c_k(n-k)\Omega^{n-k-1}\omega^k > 0$. Let ϕ satisfying $\sup_X \phi = 0$ be a smooth solution of*

$$(2.1) \quad \Omega_\phi^n = \sum_{k=0}^{n-1} \binom{n}{k} c_k \Omega_\phi^{n-k} \omega^k$$

where $c_k \geq 0$ are smooth functions. Then $\|\phi\|_{C^0} \leq C(c_k, \Omega, \omega, X)$.

The proof of this lemma follows almost word-to-word from [10] and is hence omitted. Alternatively one can use the “standard” Moser iteration type argument ([9] for instance).

Next we prove a Laplacian bound for generalised Monge-Ampère equations with “non-constant coefficients” (an improvement of a result of Sun [9] in the Kähler case). Using Evans-Krylov theory one can prove that $\|\phi\|_{C^{2,\alpha}} \leq C$. We change the normalisation of ϕ to be $\inf \phi = 1$. For this estimate we need the uniform estimate.

Proof of theorem 1.1 :

Define $\sigma_k(A)$ as the coefficient of t^k in $\det(I + tA)$ and $S_k(A) = \frac{\sigma_{n-k}(A)}{\sigma_n(A)}$ for Hermitian matrices A whose eigenvalues will be denoted by λ_i . Recall that for diagonal positive matrices A and arbitrary Hermitian matrices B we have the following standard result (for instance lemma 8 in [2]). We denote all constants by C in whatever follows.

$$(2.2) \quad \sum \frac{\partial^2 S_k}{\partial a_{ij} \partial a_{\bar{k}l}} B_{ij} \bar{B}_{kl} + \sum \frac{\partial S_k}{\partial \lambda_i} \frac{|B_{ij}|^2}{\lambda_j} \geq 0.$$

Akin to Yau's celebrated proof of the Calabi conjecture, let $\psi = e^{-\gamma(\phi)} \frac{n\omega^{n-1}\Omega_\phi}{\omega^n}$ be a function where $\gamma(x)$ is to be chosen later. Assume that the maximum of ψ is attained at p and that coordinates are chosen near p so that $\omega(p) = \sqrt{-1} \sum dz^i \wedge d\bar{z}^i$, $\omega_{,i}(p) = \omega_{,\bar{i}}(p) = 0$, and $\Omega_\phi = \sqrt{-1} \sum \lambda_i dz^i \wedge d\bar{z}^i$. At p , $\nabla\psi = 0$ and $\nabla^2\psi$ is negative-definite. Using this we have,

$$\begin{aligned}
0 &= \psi_{,i}(p) = \frac{n\omega^{n-1}\Omega_{\phi,i}}{\omega^n} - \gamma' \phi_{,i} \frac{n\omega^{n-1}\Omega_\phi}{\omega^n} \\
0 &\geq \sum_{k,i} (-c_k \frac{\partial S_k}{\partial \lambda_i}) \psi_{,\bar{i}\bar{i}}(p) \\
\Rightarrow 0 &\geq \sum_{k,i} (-c_k \frac{\partial S_k}{\partial \lambda_i}) \left[\frac{n\omega^{n-1}\Omega_{\phi,i}}{\omega^n} - \gamma' \phi_{,i} \frac{n\omega^{n-1}\Omega_\phi}{\omega^n} \right] (-\gamma') \phi_{,\bar{i}} \\
&+ \sum_{k,i} (-c_k \frac{\partial S_k}{\partial \lambda_i}) \left[\frac{n(n-1)\omega^{n-2}\omega_{,\bar{i}\bar{i}}\Omega_\phi}{\omega^n} - \frac{n\omega^{n-1}\Omega_\phi}{\omega^n} \frac{n\omega^{n-1}\omega_{,\bar{i}\bar{i}}}{\omega^n} + \frac{n\omega^{n-1}\Omega_{\phi,\bar{i}\bar{i}}}{\omega^n} \right. \\
(2.3) \quad &\left. - \gamma'' |\phi_{,i}|^2 \frac{n\omega^{n-1}\Omega_\phi}{\omega^n} - \gamma' \phi_{,\bar{i}\bar{i}} \frac{n\omega^{n-1}\Omega_\phi}{\omega^n} - \gamma' \phi_{,i} \frac{n\omega^{n-1}\Omega_{\phi,\bar{i}\bar{i}}}{\omega^n} \right].
\end{aligned}$$

At this point we divide equation 1.2 on both sides by Ω_ϕ^n and differentiate once to obtain at p

$$(2.4) \quad 0 = \sum_k c_{k,\mu} S_k + \sum_k c_k \frac{\partial S_k}{\partial \lambda_i} (\Omega_\phi)_{\bar{i}\bar{i},\mu}.$$

Differentiating again and using inequality 2.2 we get

$$\begin{aligned}
0 &\geq \sum c_{k,\mu} S_k + \sum c_{k,\mu} \frac{\partial S_k}{\partial \lambda_i} (\Omega_\phi)_{\bar{i}\bar{i},\mu} + \sum c_{k,\bar{\mu}} \frac{\partial S_k}{\partial \lambda_i} (\Omega_\phi)_{\bar{i}\bar{i},\mu} + \sum c_k \binom{n}{k} \frac{\Omega_\phi^{n-k} k \omega^{k-1} \omega_{,\mu\bar{\mu}}}{\Omega_\phi^n} \\
(2.5) \quad &+ \sum c_k \frac{\partial S_k}{\partial \lambda_i} (\Omega_\phi)_{\bar{i}\bar{i},\mu\bar{\mu}} - \sum_k c_k \frac{\partial S_k}{\partial \lambda_i} \frac{|\Omega_\phi|_{i\bar{j},\mu}^2}{\lambda_j}.
\end{aligned}$$

Using 2.5 in 2.3 we get (after simplifying)

$$\begin{aligned}
C &\geq \sum_{k,i} (-c_k \frac{\partial S_k}{\partial \lambda_i}) \left[\frac{n\omega^{n-1}\Omega_{\phi,i}}{\omega^n} (-\gamma' \phi_{,\bar{i}}) + \frac{n\omega^{n-1}\Omega_{\phi,\bar{i}}}{\omega^n} (-\gamma' \phi_{,i}) + ((\gamma')^2 - \gamma'') |\phi_{,i}|^2 \frac{n\omega^{n-1}\Omega_\phi}{\omega^n} \right. \\
&\quad \left. - \gamma' \phi_{,\bar{i}\bar{i}} \frac{n\omega^{n-1}\Omega_\phi}{\omega^n} - C \frac{n\omega^{n-1}\Omega_\phi}{\omega^n} \right] \\
(2.6) \quad &+ \sum c_{k,\mu} \frac{\partial S_k}{\partial \lambda_i} (\Omega_\phi)_{\bar{i}\bar{i},\mu} + \sum c_{k,\bar{\mu}} \frac{\partial S_k}{\partial \lambda_i} (\Omega_\phi)_{\bar{i}\bar{i},\mu} - \sum_k c_k \frac{\partial S_k}{\partial \lambda_i} \frac{|\Omega_\phi|_{i\bar{j},\mu}^2}{\lambda_j}.
\end{aligned}$$

At this juncture, we notice that (here is where we use the assumption that either c_k is identically zero or $c_k > 0$ everywhere)

$$\begin{aligned}
 \sum c_{k,\bar{\mu}} \frac{\partial S_k}{\partial \lambda_i} (\Omega_\phi)_{\bar{i}\bar{i},\mu} + \sum c_{k,\mu} \frac{\partial S_k}{\partial \lambda_i} (\Omega_\phi)_{\bar{i}\bar{i},\bar{\mu}} &\geq \sum c_k \frac{\partial S_k}{\partial \lambda_i} \left(\epsilon \frac{|(\Omega_\phi)_{\bar{i}\bar{i},\mu}|^2}{\lambda_i} + \frac{1}{\epsilon} |(\ln c_k)_{,\bar{\mu}}|^2 \lambda_i \right) \\
 &= \sum c_k \frac{\partial S_k}{\partial \lambda_i} \epsilon \frac{|(\Omega_\phi)_{\bar{i}\bar{i},\mu}|^2}{\lambda_i} + \sum \frac{1}{\epsilon} |(\ln c_k)_{,\bar{\mu}}|^2 c_k k S_k \\
 &\geq \sum c_k \frac{\partial S_k}{\partial \lambda_i} \epsilon \frac{|(\Omega_\phi)_{\bar{i}\bar{i},\mu}|^2}{\lambda_i} - \frac{C}{\epsilon}.
 \end{aligned}
 \tag{2.7}$$

Likewise, using the Cauchy-Schwartz inequality on the first two terms of 2.6 we see that

$$\frac{n\omega^{n-1}\Omega_{\phi,i}}{\omega^n} (-\gamma' \phi_{,\bar{i}}) + \frac{n\omega^{n-1}\Omega_{\phi,\bar{i}}}{\omega^n} (-\gamma' \phi_{,i}) \geq -\frac{\omega^n}{n\omega^{n-1}\Omega_\phi} \left| \frac{n\omega^{n-1}\Omega_{\phi,i}}{\omega^n} \right|^2 - (\gamma')^2 |\phi_{,i}|^2 \frac{n\omega^{n-1}\Omega_\phi}{\omega^n}.
 \tag{2.8}$$

Putting 2.6, 2.7, and 2.8 together we obtain

$$\begin{aligned}
 C + \frac{C}{\epsilon} &\geq \sum_{k,i} (-c_k \frac{\partial S_k}{\partial \lambda_i}) \left[-\frac{\omega^n}{n\omega^{n-1}\Omega_\phi} \left| \frac{n\omega^{n-1}\Omega_{\phi,i}}{\omega^n} \right|^2 - (\gamma'' + C) |\phi_{,i}|^2 \frac{n\omega^{n-1}\Omega_\phi}{\omega^n} - \gamma' \phi_{,\bar{i}\bar{i}} \frac{n\omega^{n-1}\Omega_\phi}{\omega^n} \right] \\
 &\quad + \sum c_k \frac{\partial S_k}{\partial \lambda_i} \epsilon \frac{|(\Omega_\phi)_{\bar{i}\bar{i},\mu}|^2}{\lambda_i} - \sum c_k \frac{\partial S_k}{\partial \lambda_i} \frac{|(\Omega_\phi)_{i\bar{j},\mu}|^2}{\lambda_j}.
 \end{aligned}
 \tag{2.9}$$

It is easy to see (for instance using Lagrange multipliers for the second inequality) that

$$\begin{aligned}
 \sum c_k \frac{\partial S_k}{\partial \lambda_i} \frac{|(\Omega_\phi)_{\bar{i}\bar{i},\mu}|^2}{\lambda_i} - \sum c_k \frac{\partial S_k}{\partial \lambda_i} \frac{|(\Omega_\phi)_{i\bar{j},\mu}|^2}{\lambda_j} &\geq 0 \\
 -\frac{\omega^n}{n\omega^{n-1}\Omega_\phi} \left| \frac{n\omega^{n-1}\Omega_{\phi,i}}{\omega^n} \right|^2 + \sum_{j,\mu} \frac{|(\Omega_\phi)_{i\bar{j},\mu}|^2}{\lambda_j} &\geq 0.
 \end{aligned}
 \tag{2.10}$$

Using 2.10 in 2.9 we see that

$$C + \frac{C}{\epsilon} \geq \sum_{k,i} (-c_k \frac{\partial S_k}{\partial \lambda_i}) \left[-\epsilon \frac{\omega^n}{n\omega^{n-1}\Omega_\phi} \left| \frac{n\omega^{n-1}\Omega_{\phi,i}}{\omega^n} \right|^2 - (\gamma'' |\phi_{,i}|^2 + C) \frac{n\omega^{n-1}\Omega_\phi}{\omega^n} - \gamma' \phi_{,\bar{i}\bar{i}} \frac{n\omega^{n-1}\Omega_\phi}{\omega^n} \right].
 \tag{2.11}$$

Using the first equation in 2.3 we see that

$$C + \frac{C}{\epsilon} \geq \sum_{k,i} (-c_k \frac{\partial S_k}{\partial \lambda_i}) \left[-((\epsilon(\gamma')^2 + \gamma'')) |\phi_{,i}|^2 + C) \frac{n\omega^{n-1}\Omega_\phi}{\omega^n} - \gamma' \phi_{,\bar{i}\bar{i}} \frac{n\omega^{n-1}\Omega_\phi}{\omega^n} \right].
 \tag{2.12}$$

Using another standard lemma (originally due to Fang-Lai-Ma [4]) in the form of lemma 4.1 in [9] we see that (assuming that $\gamma' \geq 0$)

$$C + \frac{C}{\epsilon} \geq \sum_{k,i} (-c_k \frac{\partial S_k}{\partial \lambda_i}) \left[-((\epsilon(\gamma')^2 + \gamma'')) |\phi_{,i}|^2 + C) \frac{n\omega^{n-1}\Omega_\phi}{\omega^n} \right] + \theta \gamma' \frac{n\omega^{n-1}\Omega_\phi}{\omega^n} \left[1 + \sum -c_k \frac{\partial S_k}{\partial \lambda_i} \right],
 \tag{2.13}$$

for some uniform constant $\theta > 0$. Choosing $\gamma(x) = \frac{\ln(x)}{\epsilon}$ we see that $-\gamma'' - \epsilon(\gamma')^2 = 0$, and $\gamma' \geq \frac{1}{\epsilon \sup \phi}$.

Choosing ϵ to be sufficiently small we see that $\frac{n\omega^{n-1}\Omega_\phi}{\omega^n}(p)$ has to be bounded above to avoid a contradiction. $\square$

Proof of theorem 1.2 : Using the method of continuity as in [8] one can (re)prove an existence result for the generalised Monge-Ampère PDE in the case where c_k are constant [8, 9, 2]. We rewrite equation 1.1 as follows.

If $n = 2m + 1$, let $\Omega = \sqrt{-1}F_0 - \omega \tan(\hat{\theta})$.

$$\begin{aligned}
\text{Im}((\omega + \sqrt{-1}(\Omega_\phi + \omega \tan(\hat{\theta})))^{2m+1}) &= \tan(\hat{\theta}) \text{Re}((\omega + \sqrt{-1}(\Omega_\phi + \omega \tan(\hat{\theta})))^{2m+1}) \\
&\Rightarrow \sum_{k=0}^{2m+1} \binom{2m+1}{k} \omega^{2m+1-k} \Omega_\phi^k \text{Im}((\sqrt{-1})^k (1 + \sqrt{-1} \tan(\hat{\theta}))^{2m+1-k}) \\
&= \tan(\hat{\theta}) \sum_{k=0}^{2m+1} \binom{2m+1}{k} \omega^{2m+1-k} \Omega_\phi^k \text{Re}((\sqrt{-1})^k (1 + \sqrt{-1} \tan(\hat{\theta}))^{2m+1-k}) \\
&\Rightarrow \sum_{k=0}^{2m+1} \binom{2m+1}{k} \sec(\hat{\theta})^{2m+1-k} \omega^{2m+1-k} \Omega_\phi^k \text{Im}((\sqrt{-1})^k (\cos((2m+1-k)\hat{\theta}) + \sqrt{-1} \sin((2m+1-k)\hat{\theta}))) \\
(2.14) \quad &= \tan(\hat{\theta}) \sum_{k=0}^{2m+1} \binom{2m+1}{k} \sec(\hat{\theta})^{2m+1-k} \omega^{2m+1-k} \Omega_\phi^k \text{Re}((\sqrt{-1})^k (\cos((2m+1-k)\hat{\theta}) + \sqrt{-1} \sin((2m+1-k)\hat{\theta})))
\end{aligned}$$

If $n = 2m$, let $\Omega = \sqrt{-1}F_0 + \omega \cot(\hat{\theta})$. Calculations similar to 2.14 show that

$$\begin{aligned}
&\cot(\hat{\theta}) \sum_{k=0}^{2m} (-1)^k \binom{2m}{k} \csc(\hat{\theta})^{2m-k} \omega^{2m-k} \Omega_\phi^k \sin((2m-k)\hat{\theta}) \\
(2.15) \quad &= \sum_{k=0}^{2m} (-1)^k \binom{2m}{k} \csc(\hat{\theta})^{2m-k} \omega^{2m-k} \Omega_\phi^k \cos((2m-k)\hat{\theta})
\end{aligned}$$

In either case, 1.1 boils down to

$$(2.16) \quad \Omega_\phi^n = \sum c_k \Omega_\phi^k \omega^{n-k}$$

where if $n = 2m + 1$

$$\begin{aligned}
c_k &= (-1)^m \binom{2m+1}{k} \sec(\hat{\theta})^{2m+1-k} \left(\tan(\hat{\theta}) \text{Re}((\sqrt{-1})^k (\cos((2m+1-k)\hat{\theta}) + \sqrt{-1} \sin((2m+1-k)\hat{\theta}))) \right. \\
(2.17) \quad &\quad \left. - \text{Im}((\sqrt{-1})^k (\cos((2m+1-k)\hat{\theta}) + \sqrt{-1} \sin((2m+1-k)\hat{\theta}))) \right),
\end{aligned}$$

and if $n = 2m$

$$(2.18) \quad c_k = \csc(\hat{\theta})^{2m-k} (-1)^k \binom{2m}{k} \left[\cot(\hat{\theta}) \sin((2m-k)\hat{\theta}) - \cos((2m-k)\hat{\theta}) \right].$$

Using the assumptions of the theorem and the comments in the beginning of the proof we obtain the required *a priori* estimates for non-constant phase and existence result for constant phase.

3. TORIC VARIETIES

We first state and prove a theorem that is almost the same as theorem 1.5 with one extremely important difference, namely, c_1 is assumed to be positive. The proof of the following theorem follows [2] very closely (we just use more general $C^{2,\alpha}$ estimates borrowed from [7]).

Theorem 3.1. *Assuming the same hypotheses as in theorem 1.5 and that $c_1 > 0$ we have the same conclusion as in theorem 1.5, i.e., the generalised Monge-Ampère equation has a solution.*

Proof of theorem 3.1 : Just as in [2] the proof proceeds by induction on the dimension n of M . For one dimensional manifolds the result is straightforward. Consider the method of continuity

$$t = \sum_{k=1}^n c_k \binom{n}{k} \frac{\omega^k \Omega_{\phi_t}^{n-k}}{\Omega_{\phi_t}^n} + d_t \frac{\omega^n}{\Omega_{\phi_t}^n}.$$

For $t \rightarrow \infty$ there exists a solution using the Calabi-Yau theorem and the implicit function theorem. Thus the problem reduces to proving that the infimum of $t \geq 0$ for which the equation has a solution is 0. For this one needs $C^{2,\alpha}$ a priori estimates on ϕ . Using the induction hypothesis one sees that the equation

$$t \Omega_{\phi_i}^{n-1} = \sum_{k=1}^{n-1} c_k \binom{n-1}{k} \omega^k \Omega_{\phi_i}^{n-1-k} + d_i \omega^{n-1}$$

has a smooth solution over each of the toric divisors D_i of X . Using the ϕ_i , just as in [2] one can construct a viscosity supersolution ϕ to the equation in a neighbourhood U of the union of the toric divisors $D = \cup_i D_i$ and use it to reduce the C^2 estimate on U to a C^2 estimate on ∂U . If one manages to prove a $C^{2,\alpha}$ estimate outside U , then one can use a lemma of Tosatti and Weinkove [11] to prove a global C^2 estimate. The Evans-Krylov theorem then gives $C^{2,\alpha}$ estimates thus completing the proof.

In order to prove $C^{2,\alpha}$ estimates outside U , one just needs to prove such estimates on every compact set K outside U . Since we are dealing with torus invariant metrics, we may assume that locally $\Omega = \sqrt{-1} \partial \bar{\partial} g$ and $\omega = \sqrt{-1} \partial \bar{\partial} f$ where g and f are convex. By using the moment map (i.e. ∇g) one can easily see find a C^1 estimate (lemma 26 in [2]). Given a C^2 estimate, one can obtain a $C^{2,\alpha}$ estimate by the Evans-Krylov theorem.

To prove a C^2 estimate, one uses a contradiction argument in conjunction with the Legendre transform. Very roughly speaking, the Legendre transform converts $D^2 g$ to $D^2 h = (D^2 g)^{-1}$. Therefore, an upper bound on $D^2 g$ translates into a lower bound on $D^2 h$. If this is violated and $D^2 h$ becomes degenerate somewhere, then a theorem of Bian-Guan ensures that it is degenerate everywhere (this is one of the places where we use the hypothesis that $c_1 > 1$) and this provides the desired contradiction. The details of this idea are in proposition 27 of [2]. However, in order to use the contradiction argument we need to take a limit (it is to that limiting h that we apply the Bian-Guan theorem). To this end, one needs the following lemma (essentially a $C^{2,\alpha}$ estimate) :

Lemma 3.2. *Suppose that $h : B \rightarrow \mathbb{R}$ is a smooth convex function on the unit ball in $\mathbb{R}^n$ satisfying*

$$(3.1) \quad \sum_{k=2}^n c_k S_{k, b_k(\nabla h)}(h_{ij}) + \sum_{i,j} a_{ij} (\nabla h) h_{ij} = 1,$$

where b_k are positive definite matrix-valued functions, $c_i \geq 0$ are constants, $0 < \lambda < a_{ij} < \Lambda$, $a_{ij}, b_k \in C^{1,\alpha}$, and $S_{k, b_k(\nabla h)}(h_{ij})$ is the coefficient of t^k in $\det(b_k(\nabla h(x)) + t D^2 h(x))$. Then we have $\|h\|_{C^{2,\alpha}(\frac{1}{2}B)} < C$ for a constant C depending on $\|h\|_{C^1}$, λ , Λ and $C^{1,\alpha}$ bounds for a_{ij} and b_k .

The proof of this lemma follows from the results in [7] and repeating the blowup argument almost word-to-word of proposition 29 in [2]. $\square$

Now we use a small trick to complete the proof of theorem 1.5.

Proof of theorem 1.5 : Assume that $c_1 = 0$. Since there are only finitely many toric subvarieties, the assumptions of theorem 1.5 imply that

$$\int_X \Omega^n - \int_X n\epsilon\omega\Omega^{n-1} - \sum_{k=2}^{n-1} \int_X \binom{n}{k} c_k \omega^k \Omega^{n-k} - (c_n - C\epsilon) \int_X \omega^n > 0$$

for some constant C satisfying $C > \frac{\int n\omega\Omega^{n-1}}{\int \omega^n}$ and a small constant $\epsilon < \frac{c_n}{C}$, and likewise for all p -dimensional subvarieties. Therefore, by theorem 3.1 we see that there is a smooth function u_ϵ and a constant $d_\epsilon > 0$ such that

$$(3.2) \quad \Omega_{u_\epsilon}^n = n\epsilon\omega\Omega_{u_\epsilon}^{n-1} + \sum_{k=2}^{n-1} c_k \binom{n}{k} \omega^k \Omega_{u_\epsilon}^{n-k} + d_\epsilon \omega^n.$$

This implies that Ω_{u_ϵ} satisfies

$$(3.3) \quad n\Omega_{u_\epsilon}^{n-1} - \sum_{k=2}^{n-1} c_k \binom{n}{k} (n-k) \omega^k \Omega_{u_\epsilon}^{n-k-1} > 0$$

Taking Ω_{u_ϵ} as the background metric instead of Ω , using the *a priori* estimates in lemma 1.1 (or alternatively in [2, 9]), and the method of continuity as in [8] we can solve the equation. This gives the desired result.

We are ready to prove corollary 1.6. Indeed,

Proof of corollary 1.6 : In the proof of theorem 1.2 we reduced the dHYM equation to a generalised Monge-Ampère equation (equation 2.16). We may deduce the desired result by applying theorem 1.5 provided the relevant hypotheses are satisfied. Indeed, lemma 8.2 of [1] implies that the “integration over subvarieties” hypotheses is equivalent to assumption (1) of corollary 1.6.

REFERENCES

- [1] T. Collins, A. Jacob, and S.T. Yau, (1, 1) forms with specified Lagrangian phase : A priori estimates and algebraic obstructions. arXiv : 1508.01934 (preprint).
- [2] T. Collins and G. Székelyhidi, Convergence of the J-flow on toric manifolds. arXiv : 1412.4809 (preprint).
- [3] X.X. Chen, On the lower bound of the Mabuchi energy and its application. *International Mathematical Research Notices*, 2000.12 (2000): 607-623.
- [4] F. Hao, M. Lai, X. Ma, On a class of fully nonlinear flows in Kähler geometry. *Journal für die reine und angewandte Mathematik (Crelles Journal)*, 2011.653 (2011): 189-220.
- [5] A. Jacob and S.T. Yau, A special Lagrangian type equation for holomorphic line bundles. arXiv : 1411.7457 (preprint)
- [6] C. Leung, S-T Yau, and E. Zaslow, From special Lagrangian to Hermitian-Yang-Mills via Fourier-Mukai transform. *Winter school on mirror symmetry, vector bundles, and Lagrangian submanifolds (Cambridge, MA, 1999)* : 209-225.
- [7] V. Pingali, $C^{2,\alpha}$ estimates and existence results for certain nonconcave PDE. arXiv : 1504.00963 (preprint).
- [8] V. Pingali, A priori estimates for a generalised Monge-Ampère PDE on some compact Kähler manifolds. arXiv : 1505.04358 (preprint) - To appear in *Complex Variables and elliptic equations*.
- [9] W. Sun, Generalized complex Monge-Ampère type equations on closed Hermitian manifolds. arXiv: 1412.8192.
- [10] G. Székelyhidi, Fully nonlinear elliptic equations on compact Hermitian manifolds. arXiv:1501.02762.
- [11] V. Tosatti and B. Weinkove, Estimates for the complex Monge-Ampère equation on Hermitian and balanced manifolds. *Asian journal of mathematics*, 14.1 (2010) :19-40.

DEPARTMENT OF MATHEMATICS, 412 KRIEGER HALL, JOHNS HOPKINS UNIVERSITY,, BALTIMORE, MD 21218, USA
E-mail address: vpingali@math.jhu.edu